

# Positive affect predicts cerebral glucose metabolism in late middle-aged adults

Christopher R. Nicholas,<sup>1,2,3</sup> Siobhan M. Hoscheidt,<sup>3</sup> Lindsay R. Clark,<sup>1,2,3</sup> Annie M. Racine,<sup>3</sup> Sara E. Berman,<sup>3</sup> Rebecca L. Kosciak,<sup>2</sup> N. Maritza Dowling,<sup>4</sup> Sanjay Asthana,<sup>1,2,3</sup> Bradley T. Christian,<sup>3,5</sup> Mark A. Sager,<sup>2</sup> and Sterling C. Johnson<sup>1,2,3</sup>

<sup>1</sup>Geriatric Research Education and Clinical Center, Wm. S. Middleton Veterans Hospital, Madison, WI, USA, <sup>2</sup>Wisconsin Alzheimer's Institute, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA, <sup>3</sup>Wisconsin Alzheimer's Disease Research Center, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA, <sup>4</sup>Department of Biostatistics & Research, School of Nursing, George Washington University, Washington, DC, USA, and <sup>5</sup>Department of Medical Physics, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Correspondence should be addressed to Sterling C. Johnson, J5/1M, Clinical Science Center, 600 Highland Ave, Madison, WI 53792, USA.  
E-mail: scj@medicine.wisc.edu.

## Abstract

Positive affect is associated with a number of health benefits; however, few studies have examined the relationship between positive affect and cerebral glucose metabolism, a key energy source for neuronal function and a possible index of brain health. We sought to determine if positive affect was associated with cerebral glucose metabolism in late middle-aged adults ( $n = 133$ ). Participants completed the positive affect subscale of the Center for Epidemiological Studies Depression Scale at two time points over a two-year period and underwent 18F-fluorodeoxyglucose-positron emission tomography scanning. After controlling for age, sex, perceived health status, depressive symptoms, anti-depressant use, family history of Alzheimer's disease, APOE  $\epsilon 4$  status and interval between visits, positive affect was associated with greater cerebral glucose metabolism across para-/limbic, frontal, temporal and parietal regions. Our findings provide evidence that positive affect in late midlife is associated with greater brain health in regions involved in affective processing and also known to be susceptible to early neuropathological processes. The current findings may have implications for interventions aimed at increasing positive affect to attenuate early neuropathological changes in at-risk individuals.

**Key words:** positive affect; cerebral glucose metabolism; well-being; FDG-PET; health; neuroimaging

## Introduction

Positive affect refers to pleasant emotional and mood states, broadly experienced as happiness, enthusiasm, contentment, coping, well-being and optimism (Folkman and Moskowitz, 2000; Lyubomirsky *et al.*, 2005; Pressman and Cohen, 2005). Over the past few decades research has provided evidence of the association between positive affect and beneficial outcomes

across many important psychosocial (Connolly and Viswesvaran, 2000; Fredrickson *et al.*, 2003; Kashdan and Roberts, 2004; Lyubomirsky *et al.*, 2005; Ruvolo, 1998; Wright and Staw, 1999) and health domains (Boehm and Kubzansky, 2012; Davidson *et al.*, 2010; Ostir *et al.*, 2001; Steptoe *et al.*, 2005). Furthermore, studies have demonstrated that positive affect and related measures of happiness and well-being are associated with healthy aging and reduced mortality (Danner *et al.*,

Received: 18 March 2016; Revised: 2 January 2017; Accepted: 1 March 2017

Published by Oxford University Press 2017. This work is written by US Government employees and is in the public domain in the US.

2001; Diener and Chan, 2011; Krijthe et al., 2011; Levy et al., 2002; Moskowitz, 2003; Ostir et al., 2000; Steptoe et al., 2015; but see Liu et al., 2016). While the direction of causality between positive affect and beneficial outcomes, particularly within the domains of health and longevity (Liu et al., 2016; Lyubomirsky et al., 2005), requires further evaluation, it is reasonable to postulate that positive affect may also be associated with brain health.

Cerebral glucose metabolism, indexed by positron emission tomography (PET) tracer 18F-fluorodeoxyglucose (18F-FDG), is an *in vivo* measure that is widely used as a marker of brain health and function in normal aging and disease. In normal aging, decreases in cerebral metabolism have been observed in anterior brain regions (Kalpouzos et al., 2009; Pardo et al., 2007; Petit-Taboue et al., 1998) with more pronounced disruption of metabolic processes within temporoparietal regions in neurodegenerative diseases such as Alzheimer's disease (AD) (Dukart et al., 2013; Kalpouzos et al., 2009; Yoshizawa et al., 2014). Abnormal cerebral glucose metabolism has also been consistently observed within fronto-limbic regions in depression (Drevets et al., 1997; Seminowicz et al., 2004), a disorder often characterized by low levels of positive affect and high levels of negative affect (Watson and Clark, 1997; Wichers et al., 2012).

Given the above evidence of age and disease-specific changes in cerebral glucose metabolism as well as the beneficial health outcomes associated with positive affect, it is possible that levels of positive affect will also have an association with cerebral glucose metabolism. To date, little is known about the relationship between positive affect and brain health, particularly in midlife when interventions aimed at modifying risk factors for pathological aging may be most effective. In the current study, we examined the relationship between positive affect and cerebral glucose metabolism in 133 healthy late middle-aged adults who provided ratings of positive affect twice over a two-year period and then underwent FDG-PET scanning. We hypothesized that greater positive affect would be positively associated with cerebral glucose metabolism, a marker of brain health.

## Materials and methods

### Participants

This study used data from the Wisconsin Registry for Alzheimer's Prevention (WRAP) cohort (Table 1), a longitudinal registry of > 1500 cognitively healthy adults recruited in middle-age and enriched for AD risk factors of APOE  $\epsilon$ 4 carriage and parental family history of AD (Sager et al., 2005). Determination of APOE  $\epsilon$ 4 and FH status in this sample has been previously described (Jonaitis et al., 2013; Koscik et al., 2014).

A total of 133 out of 178 asymptomatic late middle-aged adults (mean age 59.29 years,  $SD = 6.11$ , range = 46.58–71.07, 71.4% female) were included in the current analyses if they had completed a comprehensive neuropsychological battery, self-report ratings of affect as indexed by the Center for Epidemiologic Studies Depression Scale (CES-D), magnetic resonance imaging (MRI), FDG-PET; and were determined by neuropsychological evaluation and consensus review to be cognitively normal, and absent of a major psychiatric diagnosis, history of neurological disease and head trauma. Forty-five participants were not included in the analyses for the following reasons: 14 declined the FDG-PET scan, 11 were lost to follow-up, nine had CES-D measurements that followed the FDG-PET scan (see section labeled 'Design'), four were MRI ineligible, four had an abnormal FDG-PET scan and three were missing CES-D

**Table 1.** Participant characteristics

Variable	Participants (N = 133)
Age at initial CES-D	59.29 (6.11)
Age at FDG-PET scan	63.22 (5.99)
Female, %	71.4
Caucasian, %	97
Education (years)	16.97 (2.97)
Initial MMSE	29.47 (0.87)
FH positive, %	66.9
APOE $\epsilon$ 4 positive, %	40.6
Perceived health status <sup>a</sup>	3.81 (0.71)
Anti-depressant use, %	30.8
CES-D total <sup>a</sup>	5.89 (4.95)
CES-D positive affect <sup>a</sup>	10.64 (1.61)
CES-D-16 depressive symptoms <sup>a</sup>	4.52 (4.01)

<sup>a</sup>Calculated as the mean across two visits.

CES-D, Center for the Epidemiologic Studies Depression; FDG-PET, fluorodeoxyglucose positron emission tomography; FH, parental family history of Alzheimer's disease; APOE  $\epsilon$ 4, Apolipoprotein Epsilon-4; MMSE, Mini-Mental Status Examination. Non-percentage values are sample means with standard deviations in parentheses.

data. The University of Wisconsin Institutional Review Board approved all study procedures and each participant signed informed consent prior to participation.

### Design

Participants completed the CES-D on two different visits scheduled approximately two years apart (mean visit 1 to visit 2 interval = 26.63 months,  $SD = 3.47$ ) prior to the FDG-PET scan. Participants then underwent FDG-PET scanning ~20 months after their second visit (mean visit 1 to FDG-PET scan interval = 47.23 months;  $SD = 15.46$  and mean visit 2 to FDG-PET scan interval = 20.60 months;  $SD = 15.85$ ).

### Self-report measures

The CES-D, a well-validated questionnaire composed of 20-items designed to measure frequency of depressive symptoms experienced over the past week (Radloff, 1977) was used to assess affect. Factor analytic research on the internal structure of the CES-D typically yields a four-factor structure (Positive Affect, Negative Affect, Somatic Symptoms and Interpersonal Problems) (Shafer, 2006). This structure was tested and confirmed in the current study sample. The four statements comprising the positive affect subscale ('I felt that I was just as good as other people'; 'I felt hopeful about the future'; 'I was happy'; 'I enjoyed life') have previously been shown to be associated with beneficial health outcomes (Ostir 2000; Moskowitz, 2003) and were used to assess positive affect in the current study. For each item, participants were asked to use a four-point Likert scale (e.g. rarely/none, some/little, occasionally/moderate, most/all of the time; scored 0–3, respectively) to indicate how often they experienced a particular statement. Scores ranged from 0 to 12, with higher scores equating to a greater frequency and/or level of positive affect. A mean positive affect score based on their ratings at the two visits was calculated for each participant as a measure of characteristic positive affect (Dotson et al., 2009a,b). We included the remaining 16-items of the CES-D (CES-D-16) to control for depressive symptoms (Stansbury et al., 2006). Scores ranged from 0 to 48, with higher scores equating to more depressive symptoms. For each

participant, scores collected at each of the two visits were used to calculate mean depressive symptoms. To assess perceived health status at each visit participants were asked, 'How would you rate your current health?' on a five-point Likert scale ranging from 'Poor' to 'Excellent', with higher scores equating to greater perceived health. A mean score of perceived health status over the two visits was calculated and included to control for potential confounding effects of ill health on positive affect (Liu et al., 2016).

### FDG-PET imaging

Each participant underwent FDG-PET imaging using the Alzheimer's Disease Neuroimaging Initiative protocol (Jagust et al., 2010) including five mCi FDG injection with a 30-min uptake period followed by a 30 min acquisition. To reduce inter-subject variability participants abstained from food, caffeine, tobacco and medication for a minimum of 4 h prior to scanning. Post-processing involved frame-to-frame realignment and summation of the 30 min emission scans. Images were co-registered to the T1-weighted image, spatially normalized to the ICBM 152 atlas and smoothed to 8 mm using SPM12 ([www.fil.ion.ucl.ac.uk/spm](http://www.fil.ion.ucl.ac.uk/spm)). Intensity normalization was achieved by calculating the mean signal over the pons and cerebellar vermis (Jagust and Landau, 2012; Knopman et al., 2013) producing scaled images for each participant that were used in second level analyses.

### MRI

T1-weighted images were acquired for the purpose of coregistration of FDG-PET scans. Participants underwent scanning on a General Electric 3.0 Tesla Discovery MR750 MRI system with an eight-channel head coil (Excite HD Brain Coil, GE Healthcare). T1-weighted images were acquired in the axial plane with a 3D fast spoiled gradient-echo sequence using the following parameters: inversion time (TI) = 450 ms; repetition time (TR) = 8.1 ms; echo time (TE) = 3.2 ms; flip angle = 12°; acquisition matrix = 256 mm × 256 mm, field of view (FOV) = 256 mm; slice thickness = 1.0 mm.

### Family history and APOE ε4 status

To verify the diagnosis of AD in the parent, parental medical records, autopsy reports or results of the dementia questionnaire (Kawas et al., 1994) were obtained and reviewed by a multidisciplinary diagnostic consensus conference (Jonaitis et al., 2013; Kosciak et al., 2014; Sager et al., 2005). Absence of FH of AD was verified through detailed medical history surveys and phone interview with the participants. Genotyping for APOE ε4 status was done previously in WRAP and described elsewhere (Johnson et al., 2011).

### Statistical analyses

Associations between positive affect and glucose metabolism were assessed using multiple regression analyses conducted in SPM12. The model included positive affect as the main continuous predictor variable with FDG-PET as the dependent measure. Age at visit 1, sex, depressive symptoms, anti-depressant use at visit 1 and/or visit 2, perceived health status, family history of AD, APOE ε4 status, and interval between visit 1 CES-D and scanning were treated as covariates (Table 1). No covariate exceeded a Pearson product correlation of 0.5. Voxel-wise analyses were restricted to gray matter using an explicit mask made by thresholding the International Consortium for Brain Mapping

Tissue Probabilistic Atlases gray matter map at 0.20. This was done to minimize type 1 errors, ensuring that voxels with less than a 20% likelihood of being gray matter were not included in analyses. Statistical parametric maps were thresholded at  $P \leq 0.001$  (uncorrected) with a cluster extent threshold of  $> 100$  contiguous voxels. Resulting clusters were considered significant if they surpassed a threshold of  $P = 0.05$ , cluster-level Family Wise Error (FWE) correction.

## Results

Table 1 provides information on the participant characteristics and self-report measures. Whole brain voxel-wise analysis revealed a significant main effect of positive affect on cerebral glucose metabolism ( $P < 0.05$  FWE cluster-corrected) in the bilateral posterior cingulate gyrus/precuneus; left angular/supramarginal gyrus; left middle/inferior temporal gyrus and left middle/frontal gyrus where greater positive affect was associated with higher cerebral glucose metabolism in these regions (See Figure 1 and Table 2 for maxima and submaxima within each cluster).

## Discussion

The current study examined the novel association between positive affect and cerebral glucose metabolism in a healthy late middle-aged sample. Consistent with our hypothesis, whole-brain voxel-wise analyses revealed a significant association between positive affect and cerebral glucose metabolism, such that greater positive affect was positively associated with relatively greater cerebral glucose metabolism within para-/limbic, frontal, temporal and parietal regions. Broadly, these regions have been implicated in functional neuroimaging studies of affective processing (Blood et al., 1999; Damasio et al., 2000; Davidson and Irwin, 1999; Duerden et al., 2013; Hofer et al., 2007; Maddock et al., 2003; Phillips et al., 2008) and show changes in glucose metabolism in depression (Fitzgerald et al., 2008; Su et al., 2014), a disorder often characterized by low levels of positive affect (Watson and Clark, 1997; Wichers et al., 2012). In addition, regional grey matter volume in the precuneus, parahippocampus, medial prefrontal cortex and insula correlate with models of well-being (Kong et al., 2015; Lewis et al., 2014); which are constructs related to positive affect and often found to be associated with beneficial health outcomes (Boehm and Kubzansky, 2012; Ryan and Deci, 2001).

The positive association between positive affect and glucose metabolism across para-/limbic, frontal, temporal and parietal regions may be informed by a constructionist perspective, in which a variety of more 'basic psychological operations' from a complex array of distributed brain networks (Lindquist et al., 2012) are involved in the monitoring, processing and generation of affect (Damasio et al., 2000; Davidson, 2004; Lindquist et al., 2015). As part of the paralimbic cortex, the PCC/precuneus, parahippocampus and insula each play a role as integrative hubs for a variety of functional processes (Buckner et al., 2009) and are involved in affective processing (Blood et al., 1999; Damasio et al., 2000; Davidson and Irwin, 1999; Duerden et al., 2013; Maddock et al., 2003; Phillips et al., 2008). The PCC/precuneus is a highly active metabolic region (Raichle et al., 2001) involved in facilitating the integration of interoceptive and exteroceptive stimuli across a spectrum of attention (Leech and Sharp, 2014). The core underlying function of the parahippocampus is to bind contextual features (e.g. emotional content of a situation or experience) and link these features to a larger network of relevant

**Table 2.** Brain regions that showed a significant main effect of positive affect on glucose metabolism using voxel-wise analyses

Cluster location	Hemisphere	k	x	y	z	Peak t	P (FWE)
Posterior cingulate gyrus (23, 29, 31), precuneus (7)	B	1832	-6	-60	16	4.76	0.001
Angular/supramarginal gyrus (39, 40), insula (13), transverse temporal gyrus (41)	L	1320	-38	-54	30	4.63	0.003
Middle temporal gyrus (21, 22), para-hippocampal gyrus, hippocampal gyrus, inferior temporal gyrus (20)	L	738	-54	-42	-6	4.43	0.031
Middle frontal gyrus (10), superior frontal gyrus (10/11), anterior cingulate gyrus (32)	L	652	-30	50	0	4.38	0.044

X, Y, Z coordinates of the peak voxel of each cluster are presented in Montreal Neurological Institute (MNI) space. Corresponding Brodmann areas are in parentheses. B = Bilateral; L, Left; k, number of contiguous voxels in cluster.




Fig. 1. Statistical parametric maps were thresholded at an uncorrected voxelwise threshold of  $P < 0.001$  with a cluster extent threshold of  $> 100$  contiguous voxels. Resulting clusters were considered significant if they surpassed cluster-level threshold of  $P < 0.05$  with a Family Wise Error correction for multiple comparisons. Sagittal and coronal slices were derived from a normalized T1 anatomical image.

memories in the association cortex (Aminoff et al., 2013; Eichenbaum et al., 2007). The insula is positioned amidst these regions with specific connections to the parahippocampus, where it is involved in integrating internal homeostatic, visceral and sensory signals (Craig, 2009, 2010; Nagai et al., 2007; Suzuki, 2009). The frontal regions spanning the middle and superior frontal gyrus, and anterior cingulate gyrus, have extensive connections to the para-/limbic system and subcortical structures (Öngür and Price, 2000), and support range of cognitive functions relevant to affective processing including integration of sensory and visceral information, evaluation and attribution of affective stimuli and decision-making and self-regulation (Bechara et al., 2000; Ochsner and Gross, 2005; Phillips et al., 2008). Lastly, the left angular/supramarginal gyri are also involved in emotional processing (Hofer et al., 2007) and considered a heteromodal convergence zone supporting the integration and conceptualization of information from sensory, language and memory subsystems (Binder and Desai, 2011; Ciaramelli et al., 2008; Seghier, 2013). Taken together, we broadly suggest, that affect emerges when integrated representations (e.g. PCC/precuneus; angular/supramarginal gyrus; insula) of internal and external sensory stimuli based on contextual information from memories (e.g. para-/hippocampal gyrus) are deemed relevant and meaningful to the individual (e.g. prefrontal/frontal cortices). Of course, this is only a partial account based on our findings, thus evidence of additional regions (e.g. amygdala) and models of affective processing (e.g. lateralization; localist) (Davidson and Irwin, 1999; Rohr et al., 2013) deserve consideration.

Within our late middle-aged sample, the association between positive affect and cerebral glucose metabolism was observed in a number of brain regions that are known to exhibit progressive reductions in metabolism in prodromal stages of

AD, including the PCC, precuneus, para-/hippocampus and temporal/parietal (Buckner et al., 2005; Chetelat et al., 2003; Minoshima et al., 1997; Mosconi, 2005; Mosconi et al., 2009; Protas et al., 2013). Affective changes associated with depressive symptoms may be an important predictor of cognitive decline and dementia (da Silva et al., 2013; Ownby et al., 2006; Stepaniuk et al., 2008), and have been shown to correspond to alterations in brain volume and cerebral blood flow in vulnerable brain regions of older adults (Dotson et al., 2009a,b; Ries et al., 2009). Given that we controlled for depressive symptoms, the relationship between positive affect and cerebral glucose metabolism, notably in brain regions that support integrative functions involved in affective processing, suggests positive affect may be an important factor associated with brain health in late midlife, when neuropathological processes are only beginning to emerge and may still be modifiable (Jessen et al., 2014; Sperling et al., 2011).

This may have implications for interventions aimed at increasing and/or maintaining positive affect (Layous et al., 2011; Proyer et al., 2014; Seligman et al., 2005; Sin and Lyubomirsky, 2009). Specifically, the 'broaden and build' theory posits that positive affect expands the scope of 'thought-action repertoires' (e.g. exploration, creative problem solving, cognitive flexibility, broadening of attention), leading to a building of resources and beneficial outcomes (Fredrickson, 2001, 2013; Fredrickson et al., 2008). Consistent with this theory, positive affect is associated with reduced morbidity for a variety of known risk factors (e.g. cardiovascular; chronic stress) for AD (Boehm and Kubzansky, 2012; Pressman and Cohen, 2005; Steptoe et al., 2005), enhanced cognitive functioning (Ashby et al., 1999; Carpenter et al., 2013; Yang et al., 2013), and as our work suggests, relatively higher levels of cerebral glucose metabolism in regions vulnerable to early neuropathological changes. Thus,

interventions targeting positive affect could have potential in mitigating pathological processes associated with cognitive and functional decline through increasing flexible and adaptive responses (e.g. thought-action repertoires) to environmental demands and/or optimizing cerebral glucose metabolism within these particular brain regions. Of course, this is highly speculative, and warrants examination with more sophisticated modeling techniques and measures to further clarify the causal relationships between these variables.

There are potential limitations of our study that deserve mentioning. First, because over half the participants in this study have a family history of AD (66.9%), were late middle-aged ( $m = 59.29$  yrs-old), highly educated ( $m = 16.97$  yrs) and predominantly Caucasian (97.0%), the generalizability of our results is limited and would benefit from replication in more diverse samples. Second, while we included a measure of perceived health, which has been shown to attenuate the association between happiness and all-cause mortality (Liu et al., 2016), we did not examine a variety of specific psychological (e.g. personality, coping style), health (e.g. cardiovascular, endocrine, exercise) and social factors (e.g. social stress and support, income) that may be associated with positive affect (Lyubomirsky et al., 2005). Lastly, we speculate based on the literature regarding possible connectivity between regions, thus future analyses employing connectivity methods for FDG-PET data would be useful.

In conclusion, this study provides initial evidence that positive affect is associated with cerebral glucose metabolism in brain regions that support a variety of integrative functions involved in affective processing. A number of these regions are also known to be vulnerable to early progressive neurodegenerative changes, and therefore lay the foundation for future studies investigating the longitudinal relationship between positive affect and factors associated with pathological aging and brain health. Additionally, our findings highlight the potential for interventions aimed at increasing and/or maintaining positive affect, particularly in late-middle aged adults when early pre-clinical neuropathological processes may still be modifiable.

## Acknowledgements

This research was supported by NIH grant AG021155 (S.C.J.), AG027161 (S.C.J.), P50 AG033514 (S.A.), NIA T32AG000213 (S.A.); University of Wisconsin Institute for Clinical and Translational Research, funded through a National Center for Research Resources/National Institutes of Health Clinical and Translational Science Award, UL1TR000427, a program of the National Center for Research Resources, United States National Institutes of Health; and Veterans Administration Geriatrics Research and Clinical Center (GRECC) Fellowship in Advanced Geriatrics and Aging at the William S. Middleton Memorial Veterans Hospital, Madison, WI. We want to thank our dedicated participants.

**Conflict of interest.** None declared.

## References

Aminoff, E.M., Kveraga, K., Bar, M. (2013). The role of the parahippocampal cortex in cognition. *Trends in Cognitive Sciences*, *17*(8), 379–90.

Asby, F.G., Isen, A.M., Turken, A.U. (1999). A neuropsychological theory of positive affect and its influence on cognition. *Psychological Review*, *106*(3), 529–50. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/10467897>.

Bechara, A., Damasio, H., Damasio, A.R. (2000). Emotion, decision making and the orbitofrontal cortex. *Cerebral Cortex*, *10*(3), 295–307.

Binder, J.R., Desai, R.H. (2011). The neurobiology of semantic memory. *Trends in Cognitive Sciences*, *15*(11), 527–36.

Blood, A.J., Zatorre, R.J., Bermudez, P., Evans, A.C. (1999). Emotional responses to pleasant and unpleasant music correlate with activity in paralimbic brain regions. *Nature Neuroscience*, *2*(4), 382–7.

Boehm, J.K., Kubzansky, L.D. (2012). The heart's content: the association between positive psychological well-being and cardiovascular health. *Psychological Bulletin*, *138*(4), 655.

Buckner, R.L., Sepulcre, J., Talukdar, T., et al. (2009). Cortical hubs revealed by intrinsic functional connectivity: mapping, assessment of stability, and relation to Alzheimer's disease. *Journal of Neuroscience*, *29*(6), 1860–73.

Buckner, R.L., Snyder, A.Z., Shannon, B.J., et al. (2005). Molecular, structural, and functional characterization of Alzheimer's disease: evidence for a relationship between default activity, amyloid, and memory. *Journal of Neuroscience*, *25*(34), 7709–17.

Carpenter, S.M., Peters, E., Västfjäll, D., Isen, A.M. (2013). Positive feelings facilitate working memory and complex decision making among older adults. *Cognition & Emotion*, *27*(1), 184–92.

Chetelat, G., Desgranges, B., de la Sayette, V., Viader, F., Eustache, F., Baron, J.C. (2003). Mild cognitive impairment: can FDG-PET predict who is to rapidly convert to Alzheimer's disease? *Neurology*, *60*(8), 1374–7. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/12707450>.

Ciaramelli, E., Grady, C.L., Moscovitch, M. (2008). Top-down and bottom-up attention to memory: a hypothesis (AtoM) on the role of the posterior parietal cortex in memory retrieval. *Neuropsychologia*, *46*(7), 1828–51.

Connolly, J.J., Viswesvaran, C. (2000). The role of affectivity in job satisfaction: a meta-analysis. *Personality and Individual Differences*, *29*(2), 265–81.

Craig, A.D. (2009). How do you feel—now? The anterior insula and human awareness. *Nature Reviews Neuroscience*, *10*(1), 59–70.

Craig, A.D. (2010). The sentient self. *Brain Structure & Function*, *214*(5–6), 563–77.

da Silva, J., Gonçalves-Pereira, M., Xavier, M., Mukaetova-Ladinska, E.B. (2013). Affective disorders and risk of developing dementia: systematic review. *The British Journal of Psychiatry*, *202*(3), 177–86.

Damasio, A.R., Grabowski, T.J., Bechara, A., et al. (2000). Subcortical and cortical brain activity during the feeling of self-generated emotions. *Nature Neuroscience*, *3*(10), 1049–56.

Danner, D.D., Snowdon, D.A., Friesen, W.V. (2001). Positive emotions in early life and longevity: findings from the nun study. *Journal of Personality and Social Psychology*, *80*(5), 804–13. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/11374751>.

Davidson, K.W., Mostofsky, E., Whang, W. (2010). Don't worry, be happy: positive affect and reduced 10-year incident coronary heart disease: the Canadian Nova Scotia Health Survey. *European Heart Journal*, *31*(9), 1065–70.

Davidson, R.J. (2004). Well-being and affective style: neural substrates and biobehavioural correlates. *Philosophical Transactions: Biological Sciences*, *359*(1449), 1395–411. Retrieved from: <http://www.jstor.org/stable/4142143>.

Davidson, R.J., Irwin, W. (1999). The functional neuroanatomy of emotion and affective style. *Trends in Cognitive Sciences*, *3*(1), 11–21. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/10234222>.

- Diener, E., Chan, M.Y. (2011). Happy people live longer: subjective well-being contributes to health and longevity. *Applied Psychology: Health and Well-Being*, 3(1), 1–43.
- Dotson, V.M., Beason-Held, L., Kraut, M.A., Resnick, S.M. (2009a). Longitudinal study of chronic depressive symptoms and regional cerebral blood flow in older men and women. *International Journal of Geriatric Psychiatry*, 24(8), 809–19.
- Dotson, V.M., Davatzikos, C., Kraut, M.A., Resnick, S.M. (2009b). Depressive symptoms and brain volumes in older adults: a longitudinal magnetic resonance imaging study. *Journal of Psychiatry & Neuroscience*, 34(5), 367–75. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/19721847>.
- Drevets, W.C., Price, J.L., Simpson, J.R., Jr, et al. (1997). Subgenual prefrontal cortex abnormalities in mood disorders. *Nature*, 386(6627), 824–7.
- Duerden, E.G., Arsalidou, M., Lee, M., Taylor, M.J. (2013). Lateralization of affective processing in the insula. *Neuroimage*, 78, 159–75.
- Dukart, J., Kherif, F., Mueller, K. et al. (2013). Generative FDG-PET and MRI model of aging and disease progression in Alzheimer's disease. *PLOS Computational Biology*, 9(4), e1002987.
- Eichenbaum, H., Yonelinas, A.P., Ranganath, C. (2007). The medial temporal lobe and recognition memory. *Annual Review of Neuroscience*, 30, 123–52.
- Fitzgerald, P.B., Laird, A.R., Maller, J., Daskalakis, Z.J. (2008). A meta-analytic study of changes in brain activation in depression. *Human Brain Mapping*, 29(6), 683–95.
- Folkman, S., Moskowitz, J.T. (2000). Positive affect and the other side of coping. *American Psychologist*, 55(6), 647–54. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/10892207>.
- Fredrickson, B.L. (2001). The role of positive emotions in positive psychology. The broaden-and-build theory of positive emotions. *American Psychologist*, 56(3), 218–26. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11315248>.
- Fredrickson, B.L. (2013). Positive emotions broaden and build. *Advances in Experimental Social Psychology*, 47(1), 53.
- Fredrickson, B.L., Cohn, M.A., Coffey, K.A., Pek, J., Finkel, S.M. (2008). Open hearts build lives: positive emotions, induced through loving-kindness meditation, build consequential personal resources. *Journal of Personality and Social Psychology*, 95(5), 1045.
- Fredrickson, B.L., Tugade, M.M., Waugh, C.E., Larkin, G.R. (2003). What good are positive emotions in crisis? A prospective study of resilience and emotions following the terrorist attacks on the United States on September 11th, 2001. *Journal of Personality and Social Psychology*, 84(2), 365–76.
- Hofer, A., Siedentopf, C.M., Ischebeck, A., et al. (2007). Sex differences in brain activation patterns during processing of positively and negatively valenced emotional words. *Psychological Medicine*, 37(01), 109–19.
- Jagust, W.J., Bandy, D., Chen, K., et al. (2010). The Alzheimer's Disease neuroimaging initiative positron emission tomography core. *Alzheimer's & Dementia*, 6(3), 221–9.
- Jagust, W.J., Landau, S.M. (2012). Apolipoprotein E, not fibrillar  $\beta$ -amyloid, reduces cerebral glucose metabolism in normal aging. *Journal of Neuroscience*, 32(50), 18227–33.
- Jessen, F., Amariglio, R.E., van Boxtel, M., et al. (2014). A conceptual framework for research on subjective cognitive decline in preclinical Alzheimer's disease. *Alzheimer's & Dementia*, 10(6), 844–52.
- Johnson, S.C., La Rue, A., Hermann, B.P., et al. (2011). The effect of TOMM40 poly-T length on gray matter volume and cognition in middle-aged persons with APOE  $\epsilon$ 3/ $\epsilon$ 3 genotype. *Alzheimer's & Dementia: The Journal of the Alzheimer's Association*, 7(4), 456–65.
- Jonaitis, E., La Rue, A., Mueller, K.D., Kosciak, R.L., Hermann, B., Sager, M.A. (2013). Cognitive activities and cognitive performance in middle-aged adults at risk for Alzheimer's disease. *Psychology and Aging*, 28(4), 1004–14.
- Kalpouzos, G., Chetelat, G., Baron, J.C., et al. (2009). Voxel-based mapping of brain gray matter volume and glucose metabolism profiles in normal aging. *Neurobiology of Aging*, 30(1), 112–24.
- Kashdan, T.B., Roberts, J.E. (2004). Trait and state curiosity in the genesis of intimacy: differentiation from related constructs. *Journal of Social and Clinical Psychology*, 23(6), 792.
- Kawas, C., Segal, J., Stewart, W.F., Corrada, M., Thal, L.J. (1994). A validation study of the dementia questionnaire. *Archives of Neurology*, 51(9), 901–6.
- Knopman, D.S., Jack, C.R., Wiste, H.J., et al. (2013). Selective worsening of brain injury biomarker abnormalities in cognitively normal elderly with  $\beta$ -amyloidosis. *JAMA Neurology*, 70(8), 1030–8. doi:10.1001/jamaneurol.2013.182.
- Kong, F., Ding, K., Yang, Z., et al. (2015). Examining gray matter structures associated with individual differences in global life satisfaction in a large sample of young adults. *Social Cognitive and Affective Neuroscience*, 10(7), 952–60.
- Kosciak, R.L., La Rue, A., Jonaitis, E.M., et al. (2014). Emergence of mild cognitive impairment in late middle-aged adults in the Wisconsin Registry for Alzheimer's prevention. *Dementia and Geriatric Cognitive Disorders*, 38(1–2), 16–30.
- Krijthe, B.P., Walter, S., Newson, R.S., Hofman, A., Hunink, M.G., Tiemeier, H. (2011). Is positive affect associated with survival? A population-based study of elderly persons. *American Journal of Epidemiology*, 173(11), 1298–307.
- Layouts, K., Chancellor, J., Lyubomirsky, S., Wang, L., Doraiswamy, P.M. (2011). Delivering happiness: translating positive psychology intervention research for treating major and minor depressive disorders. *The Journal of Alternative and Complementary Medicine*, 17(8), 675–83.
- Leech, R., Sharp, D.J. (2014). The role of the posterior cingulate cortex in cognition and disease. *Brain*, 137(Pt 1), 12–32.
- Levy, B.R., Slade, M.D., Kunkel, S.R., Kasl, S.V. (2002). Longevity increased by positive self-perceptions of aging. *Journal of Personality and Social Psychology*, 83(2), 261–70. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/12150226>.
- Lewis, G.J., Kanai, R., Rees, G., Bates, T.C. (2014). Neural correlates of the 'good life': eudaimonic well-being is associated with insular cortex volume. *Social Cognitive and Affective Neuroscience*, 9(5), 615–8.
- Lindquist, K.A., Satpute, A.B., Wager, T.D., Weber, J., Barrett, L.F. (2015). The brain basis of positive and negative affect: evidence from a meta-analysis of the human neuroimaging literature. *Cerebral Cortex*, 5, 1910–22.
- Lindquist, K.A., Wager, T.D., Kober, H., Bliss-Moreau, E., Barrett, L.F. (2012). The brain basis of emotion: a meta-analytic review. *Behavioral and Brain Sciences*, 35(03), 121–43.
- Liu, B., Floud, S., Pirie, K., et al. (2016). Does happiness itself directly affect mortality? The prospective UK Million Women Study. *Lancet*, 387, 874–81. doi:10.1016/S0140-6736(15)01087-9.
- Lyubomirsky, S., King, L., Diener, E. (2005). The benefits of frequent positive affect: does happiness lead to success? *Psychological Bulletin*, 131(6), 803–55.
- Maddock, R.J., Garrett, A.S., Buonocore, M.H. (2003). Posterior cingulate cortex activation by emotional words: fMRI evidence from a valence decision task. *Human Brain Mapping*, 18(1), 30–41.

- Minoshima, S., Giordani, B., Berent, S., Frey, K.A., Foster, N.L., Kuhl, D.E. (1997). Metabolic reduction in the posterior cingulate cortex in very early Alzheimer's disease. *Annals of Neurology*, *42*(1), 85–94.
- Mosconi, L. (2005). Brain glucose metabolism in the early and specific diagnosis of Alzheimer's disease. FDG-PET studies in MCI and AD. *European Journal of Nuclear Medicine and Molecular Imaging*, *32*(4), 486–510.
- Mosconi, L., Mistur, R., Switalski, R., et al. (2009). FDG-PET changes in brain glucose metabolism from normal cognition to pathologically verified Alzheimer's disease. *European Journal of Nuclear Medicine and Molecular Imaging*, *36*(5), 811–22.
- Moskowitz, J.T. (2003). Positive affect predicts lower risk of AIDS mortality. *Psychosomatic Medicine*, *65*(4), 620–6. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/12883113>.
- Nagai, M., Kishi, K., Kato, S. (2007). Insular cortex and neuropsychiatric disorders: a review of recent literature. *European Psychiatry*, *22*(6), 387–94.
- Ochsner, K.N., Gross, J.J. (2005). The cognitive control of emotion. *Trends in Cognitive Sciences*, *9*(5), 242–9.
- Öngür, D., Price, J. (2000). The organization of networks within the orbital and medial prefrontal cortex of rats, monkeys and humans. *Cerebral Cortex*, *10*(3), 206–19.
- Ostir, G.V., Markides, K.S., Black, S.A., Goodwin, J.S. (2000). Emotional well-being predicts subsequent functional independence and survival. *Journal of the American Geriatrics Society*, *48*(5), 473–8. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/10811538>.
- Ostir, G.V., Markides, K.S., Peek, M.K., Goodwin, J.S. (2001). The association between emotional well-being and the incidence of stroke in older adults. *Psychosomatic Medicine*, *63*(2), 210–5. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/11292267>.
- Ownby, R.L., Crocco, E., Acevedo, A., John, V., Loewenstein, D. (2006). Depression and risk for Alzheimer disease: systematic review, meta-analysis, and metaregression analysis. *Archives of General Psychiatry*, *63*(5), 530–8.
- Pardo, J.V., Lee, J.T., Sheikh, S.A., et al. (2007). Where the brain grows old: decline in anterior cingulate and medial prefrontal function with normal aging. *Neuroimage*, *35*(3), 1231–7.
- Petit-Taboue, M., Landeau, B., Desson, J., Desgranges, B., Baron, J. (1998). Effects of healthy aging on the regional cerebral metabolic rate of glucose assessed with statistical parametric mapping. *Neuroimage*, *7*(3), 176–84.
- Phillips, M.L., Ladouceur, C.D., Drevets, W.C. (2008). A neural model of voluntary and automatic emotion regulation: implications for understanding the pathophysiology and neurodevelopment of bipolar disorder. *Molecular Psychiatry*, *13*(9), 833–57.
- Pressman, S.D., Cohen, S. (2005). Does positive affect influence health? *Psychological Bulletin*, *131*(6), 925.
- Protas, H.D., Chen, K., Langbaum, J.B., et al. (2013). Posterior cingulate glucose metabolism, hippocampal glucose metabolism, and hippocampal volume in cognitively normal, late-middle-aged persons at 3 levels of genetic risk for Alzheimer disease. *JAMA Neurology*, *70*(3), 320–5.
- Proyer, R.T., Gander, F., Wellenzohn, S., Ruch, W. (2014). Positive psychology interventions in people aged 50–79 years: long-term effects of placebo-controlled online interventions on well-being and depression. *Aging & Mental Health*, *18*(8), 997–1005.
- Radloff, L.S. (1977). The CES-D scale: a self-report depression scale for research in the general population. *Applied Psychological Measurement*, *1*(3), 385–401.
- Raichle, M.E., MacLeod, A.M., Snyder, A.Z., Powers, W.J., Gusnard, D.A., Shulman, G.L. (2001). A default mode of brain function. *Proceedings of the National Academy of Sciences of the United States of America*, *98*(2), 676–82.
- Ries, M.L., Wichmann, A., Bendlin, B.B., Johnson, S.C. (2009). Posterior cingulate and lateral parietal gray matter volume in older adults with depressive symptoms. *Brain Imaging and Behavior*, *3*(3), 233–9.
- Rohr, C.S., Okon-Singer, H., Craddock, R.C., Villringer, A., Margulies, D.S. (2013). Affect and the brain's functional organization: a resting-state connectivity approach. *PLoS ONE*, *8*(7), e68015.
- Ruvolo, A.P. (1998). Marital well-being and general happiness of newlywed couples: relationships across time. *Journal of Social and Personal Relationships*, *15*(4), 470–89.
- Ryan, R.M., Deci, E.L. (2001). On happiness and human potentials: a review of research on hedonic and eudaimonic well-being. *Annual Review of Psychology*, *52* (1), 141–166.
- Sager, M.A., Hermann, B., La Rue, A. (2005). Middle-aged children of persons with Alzheimer's disease: APOE genotypes and cognitive function in the Wisconsin Registry for Alzheimer's Prevention. *Journal of Geriatric Psychiatry and Neurology*, *18*(4), 245–9.
- Seghier, M.L. (2013). The angular gyrus multiple functions and multiple subdivisions. *The Neuroscientist*, *19*(1), 43–61.
- Seligman, M.E., Steen, T.A., Park, N., Peterson, C. (2005). Positive psychology progress: empirical validation of interventions. *American Psychologist*, *60*(5), 410.
- Seminowicz, D., Mayberg, H., McIntosh, A., et al. (2004). Limbic-frontal circuitry in major depression: a path modeling meta-analysis. *Neuroimage*, *22*(1), 409–18.
- Shafer, A.B. (2006). Meta-analysis of the factor structures of four depression questionnaires: Beck, CES-D, Hamilton, and Zung. *Journal of Clinical Psychology*, *62*(1), 123–46.
- Sin, N.L., Lyubomirsky, S. (2009). Enhancing well-being and alleviating depressive symptoms with positive psychology interventions: a practice-friendly meta-analysis. *Journal of Clinical Psychology*, *65*(5), 467–87.
- Sperling, R.A., Aisen, P.S., Beckett, L.A., et al. (2011). Toward defining the preclinical stages of Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's & Dementia*, *7*(3), 280–92.
- Stansbury, J.P., Ried, L.D., Velozo, C.A. (2006). Unidimensionality and bandwidth in the Center for Epidemiologic Studies Depression (CES-D) scale. *Journal of Personality Assessment*, *86*(1), 10–22.
- Stepaniuk, J., Ritchie, L., Tuokko, H.A. (2008). Neuropsychiatric impairments as predictors of mild cognitive impairment, dementia, and Alzheimer's disease. *American Journal of Alzheimer's Disease and Other Dementias*, *23*, 326–33.
- Steptoe, A., Deaton, A., Stone, A.A. (2015). Subjective wellbeing, health, and ageing. *The Lancet*, *385*(9968), 640–8.
- Steptoe, A., Wardle, J., Marmot, M. (2005). Positive affect and health-related neuroendocrine, cardiovascular, and inflammatory processes. *Proceedings of the National Academy of Sciences of the United States of America*, *102*(18), 6508–12.
- Su, L., Cai, Y., Xu, Y., Dutt, A., Shi, S., Bramon, E. (2014). Cerebral metabolism in major depressive disorder: a voxel-based meta-analysis of positron emission tomography studies. *BMC Psychiatry*, *14*(1), 1.
- Suzuki, W. (2009). Comparative analysis of the cortical afferents, intrinsic projections and interconnections of the parahippocampal region in monkeys and rats. In: Gazzaniga, M.S., editor.

- The Cognitive Neurosciences*, pp.659–774. Cambridge, MA: MIT Press.
- Watson, D., Clark, L.A. (1997). Measurement and mismeasurement of mood: recurrent and emergent issues. *Journal of Personality Assessment*, *68*(2), 267–96.
- Wichers, M., Lothmann, C., Simons, C.J., Nicolson, N.A., Peeters, F. (2012). The dynamic interplay between negative and positive emotions in daily life predicts response to treatment in depression: a momentary assessment study. *British Journal of Clinical Psychology*, *51*(2), 206–22.
- Wright, T.A., Staw, B.M. (1999). Affect and favorable work outcomes: Two longitudinal tests of the happy-productive worker thesis. *Journal of Organizational Behavior*, *20*(1), 1–23.
- Yang, H., Yang, S., Isen, A.M. (2013). Positive affect improves working memory: implications for controlled cognitive processing. *Cognition & Emotion*, *27*(3), 474–82.
- Yoshizawa, H., Gazes, Y., Stern, Y., Miyata, Y., Uchiyama, S. (2014). Characterizing the normative profile of 18F-FDG PET brain imaging: sex difference, aging effect, and cognitive reserve. *Psychiatry Research*, *221*(1), 78–85.